

TeVPA – Columbus, Ohio, USA – 7<sup>th</sup> August 2017



## NEUTRINO ASTRONOMY MEASURING THE SUN'S CORE

Based on J.H.Davis, Phys.Rev.Lett. 117 (2016) 211101

JONATHAN DAVIS (KING'S COLLEGE LONDON) JONATHAN.H.M.DAVIS@GMAIL.COM

#### **NEUTRINO ASTRONOMY**

- In order to do astronomy with neutrinos we need to be able to work out where they are coming from in the sky.
- Interested in two cases: supernova neutrinos and solar neutrinos.
- If a supernova goes off in our galaxy and we detect its neutrinos, can we work out where in the galaxy it happened?
- Can we use neutrino directionality to probe the sun's core?



### **DETECTION OF NEUTRINOS**

- I will focus on water Cherenkov detectors such as Super Kamiokande (SK).
- Very large detectors such as SK mean that it is possible to obtain large statistics even though neutrinos interact weakly.
- They are good for solar and supernova neutrinos and they have directional sensitivity for incoming neutrinos.



#### **DETECTION OF SOLAR NEUTRINOS**

- An MeV-energy neutrino scatters elastically off an electron.
- The electron emits Cherenkov light, which is observed by photomultiplier tubes.

 $\frac{\nu + e^-}{\ln H_2 O} \rightarrow \nu + e^-_{\text{Emits C}}$ **Emits Cherenkov** Light



#### **DETECTION OF NEUTRINOS**

- The elastic scattering cross section is **strongly forward-peaked** for MeV-scale neutrinos, especially for higher recoil energies.
- Hence the electron after scattering will point back towards the original direction of the neutrino.



Ando and Sato, arXiv:hep-ph/0110187v4

# THE ANGULAR RESOLUTION OF NEUTRINO DETECTORS

- Unfortunately the actual resolution is much worse, since the electron scatters in the water multiple times. Hence the observed electron direction is only weakly correlated with the incident neutrino direction.
- This multiple scattering contributes almost all of the angular resolution, and is **well-understood due to calibration data**.
- See e.g. Calibration of Super-Kamiokande using an electron LINAC, Nuclear Instruments and Methods in Physics Research A 421 (1999) 113-129.



R. Tomas et al., Phys. Rev. D68 (2003) 093013

#### CASE STUDY: SUPERNOVA POINTING WITH NEUTRINOS

- Since the multiple-scattering of the electron is well-known, we can reconstruct the supernova direction.
- Obtain a simple approximate formula for the angular resolution for SN pointing:

$$\delta\theta \approx \frac{25^{\circ}}{\sqrt{N_s}}$$



FIG. 4: Angular distribution of  $\bar{\nu}_e p \rightarrow ne^+$  events (green) and elastic scattering events  $\nu e^- \rightarrow \nu e^-$  (blue) of one simulated SN.

R. Tomas et al., Phys.Rev. D68 (2003) 093013 S. Ando and K. Sato, Prog. Theor. Phys. 107 (2002) 957 J. Beacom and P. Vogel, Phys.Rev. D60 (1999) 033007

## **SOLAR NEUTRINOS**

- Solar neutrinos are produced via fusion reactions occurring in the Sun's core.
- The solar core has a radius of 20% to 25% of the solar radius.
- Their energies and fluxes depend on the fusion reactions in which they are created.



#### WHERE ARE 8B NEUTRINOS PRODUCED IN THE CORE?

- Different fusion reactions should occur at different positions within the core.
- We focus on 8B neutrinos, which are predicted to be produced in a spherical region located at 5% of the solar radius from the core centre.



#### MAXIMUM LIKELIHOOD ANALYSIS – Generating signal distributions

- We need to generate the distribution of electrons in a water Cherenkov detector, given an assumption on the neutrino distribution in the solar core.
- Start by generating initial angles of the neutrinos as they arrive at Earth, given a distribution in the core:



#### MAXIMUM LIKELIHOOD ANALYSIS – Generating signal distributions

- Combine the initial distribution of neutrinos, the differential cross section of electron-nu scattering and the distribution of electron multi-scattering in the detector.
- Repeat this process for different initial neutrino distributions within the solar core.



#### MAXIMUM LIKELIHOOD ANALYSIS – GENERATING SIGNAL DISTRIBUTIONS



#### **MAXIMUM LIKELIHOOD ANALYSIS - RESULTS**



J. H. Davis, Phys.Rev.Lett. 117 (2016) 211101

### CONCLUSION

- We can use solar neutrino experiments as telescopes of the solar interior.
- Super Kamiokande has 20 years of data so can already constrain the solar neutrino production region to be within the solar core.
- A 500 kton experiment, perhaps Hyper Kamiokande, could do much better, but it would need to keep background levels as small or smaller than for Super Kamiokande.











Jonathan.h.m.davis@gmail.com Jonathan.davis@kcl.ac.uk