

Multi-PeV Signals from a New Astrophysical Neutrino Flux Beyond the Glashow Resonance


Ranjan Laha

Kavli Institute for Particle Astrophysics and Cosmology (KIPAC)
Stanford University and SLAC National Accelerator Laboratory



Kistler and Laha arXiv: 1605.08781 (submitted)

Discovery of astrophysical neutrinos



High Energy Starting
Event analysis --- 6 year
data set

Fit performed including
all events between
 $60 \text{ TeV} < E_{\text{dep}} < 3 \text{ PeV}$

No Glashow resonance
event


$$E^2 \phi(E) = 2.46 \pm 0.8 \times 10^{-8} \left(\frac{E}{100 \text{ TeV}} \right)^{-0.92} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$$

Steep spectral index from the HESE excess

No statistically significant clustering in space

See talks by Halzen, Kowalski, Wandkowsky, Niederhausen, Richman, Lu, Yuan, Mancina, Wood, Weiler, Ando, Murase, Petropoulou, Sudoh, Denton, Bustamante, Auguelles, Kheirandish, Stecker, Lai, Klop, Yuan, Cerruti, Rodrigues, Fang

Discovery of astrophysical neutrinos



Northern Hemisphere
through going muons
--- 6 year data set

Fit of through going
events between
 $194 \text{ TeV} < E_\nu < 7.8 \text{ PeV}$


$$\Phi_{\nu+\bar{\nu}} = 0.9_{-0.27}^{+0.30} \times 10^{18} \left(\frac{E_\nu}{100 \text{ TeV}} \right)^{-2.13 \pm 0.13} \text{GeV}^{-1} \text{cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$$

No statistically significant clustering in space

See talks by Weiler, Ando, Murase, Petropoulou, Sudoh, Denton, Bustamante, Auguelles, Kheirandish, Stecker, Lai, Klop, Yuan, Cerruti, Rodrigues, Fang

2.6 PeV track event

IceCube 1607.08006



Deposited energy 2.6 ± 0.3 PeV

Reconstructed equatorial
coordinates: decl. 11.42°
RA 110.63°

Does not point towards any
known astrophysical source

Highest energy track event
detected till date --- very
important to analyze it
thoroughly



Immediate questions:

1. What **flavor of neutrino** produces such a track?
2. What are **implications for astrophysical neutrinos** in light of prior discoveries?


What neutrino flavor produces a track?

- **Muons** are assumed to give rise to **all** through-going track like events
- To deposit 2.6 PeV of energy, the muon typically requires $\gtrsim 5$ PeV **energy at detector entry point** --- it is probable that this is a **super-Glashow** (energy ≥ 6.3 PeV) neutrino
- An overlooked possibility in the literature: **very high energy through going taus can also give rise to track-like events**
- To deposit 2.6 PeV of energy, the tau requires $\gtrsim 50$ PeV **energy at detector entry point**
- Can IceCube individually **distinguish** a **through going tau** from a **through going muon**?
- We discuss astrophysical scenarios for each of these possibilities

Muon tracks



Wandkowsky TeVPA 207





Many muon tracks observed in IceCube

Many theoretical studies on muon tracks


Muon tracks in IceCube have been well studied

Double bang v/s tau tracks

Tau tracks are produced by ν_τ with energy $\gtrsim 50$ PeV



Wandkowsky TeVPA 2017



$$\nu_\tau + N \rightarrow \tau + X$$

?

How do **tau tracks** look in IceCube?

A generic equation for event numbers for through going leptons

$$\frac{d}{dE_\ell} \left[b_\ell(E_\ell) \frac{dN_\ell}{dE_\ell} \right] + \frac{m_\ell}{c \tau_\ell E_\ell} \frac{dN_\ell}{dE_\ell} = Q(E_\ell)$$

Kistler and Laha arXiv: 1605.08781

Lepton energy loss: $b_\ell \equiv dE_\ell/dX$

Lepton energy E_ℓ

Lepton lifetime τ_ℓ


includes
Earth absorption

Volumetric source term $Q(E_\ell) \approx N_A \rho \phi_\ell \left(\frac{E_\ell}{\langle 1-y \rangle} \right) \sigma_{\text{CC}} \left(\frac{E_\ell}{\langle 1-y \rangle} \right) / \langle 1-y \rangle$

Regeneration is important for taus

Earth absorption is important for all neutrino flavors


Where does it come from?



Shaded: Sky density of $E_\mu > 5$ PeV muons from model φ_8
Solid: The horizon demarcates upgoing and downgoing directions
Dashed: The rough 10° downgoing boundary for atmospheric muons
No gamma-ray source was reported by HAWC

The 2.6 PeV track event probably comes from a diffuse astrophysical neutrino flux

Astrophysical neutrino fluxes





φ_7 and φ_8 resemble BL Lac AGN models

Rescaled combinations of φ_7 and φ_9 approximate GZK neutrinos from EBL and CMB interactions

$$\varphi_i(E_\nu) = f_i \left[\left(\frac{E_\nu}{E_i} \right)^{\alpha\eta} + \left(\frac{E_\nu}{E_i} \right)^{\beta\eta} \right]^{1/\eta} \quad \begin{matrix} i = 7, 8, \text{ and } 9 \\ E_i = 10^i \text{ GeV} \end{matrix}$$

Energy distribution



Muons of **energy $\geq 100 \text{ GeV}$** can traverse the full IceCube detector

Taus of **energy $\geq 50 \text{ PeV}$** can traverse the full IceCube detector

General conclusion: in order to deposit 2.6 PeV energy inside IceCube, the energy of tau as it enters the detector must be approximately **an order of magnitude larger** than that of muon: **enormous physical significance**

Must optimize tools for this **new signal** in IceCube

Event numbers

Kistler and Laha
arXiv: 1605.08781

Events in $5 \text{ km}^2 \text{ yr}$ (μ, τ tracks) or $5 \text{ km}^3 \text{ yr}$ (showers).

	$E_\nu^{-2.6}$	φ_7	φ_8	φ_9
upgoing μ : $E_\mu > 5 \text{ PeV}$	0.04	0.22	0.25	0.08
down μ : $E_\mu > 5 \text{ PeV}; \cos \theta_{\text{nadir}} > -0.2$	0.06	0.30	0.46	0.25
upgoing τ : $E_{\tau, \text{up}} > 50 \text{ PeV}$	—	0.01	0.08	0.07
down τ : $E_\tau > 50 \text{ PeV}; \cos \theta_{\text{nadir}} > -0.2$	—	0.03	0.17	0.19
Total tracks	0.1	0.56	0.96	0.59
$\bar{\nu}_e e$ shower: $E_{\text{em}} > 5 \text{ PeV}$	—	2.6	0.36	0.04
$\nu_e + \bar{\nu}_e$ CC: $E_{\text{em}} > 5 \text{ PeV}$	—	0.87	0.50	0.12
$\nu + \bar{\nu}$ NC: $E_{\text{em}} > 5 \text{ PeV}$	—	0.18	0.42	0.16



The normalizations of φ_7 , φ_8 , and φ_9 can be made variable
The harder spectra adopted by us is more favorable to give rise to the 2.6 PeV track event
The $E^{-2.13}$ produces too many Glashow resonance events (~ 3), and might be disfavored
Is the 2.6 PeV track event hinting at a [super-Glashow astrophysical neutrino flux?](#)

Conclusions

- Common knowledge: Muons give rise to all through going tau track events in IceCube ---- incomplete!
- We show for the first time that a through going tau can also give rise to tracks
- The through going tau track is a new signal in IceCube ---- needs more research ---- can we distinguish a through going muon track and a through going tau track ?

Angular distribution

Kistler and Laha arXiv: 1605.08781



Angular spectra of $E_\mu > 5 \text{ PeV}$ muons and $E_\tau > 50 \text{ PeV}$ taus

For the same spectra, the angular distribution is different --- through going muon tracks and through going tau tracks probe different energy ranges of the underlying astrophysical neutrino spectrum

For $\varphi_7 \rightarrow \varphi_8 \rightarrow \varphi_9$, the tau/ muon track ratio approaches unity